

A Mechanism for Ferrimagnetism and Incommensurability in One-Dimensional Systems

A. M. Tsvelik

Department of Physics, University of Oxford, 1 Keble Road, Oxford OX1 3NP, UK

In this paper I discuss a mechanism for ferrimagnetism in (1+1)-dimensions. The mechanism is related to a special class of interactions described by operators with non-zero Lorentz spin. Such operators are present in such problems as the problem of tunneling between Luttinger liquids and the problem of frustrated spin ladder. Exact solutions are presented for a representative class of models possessing a continuous isotopic symmetry. It is shown that the interactions (i) dynamically generate static oscillations with the wave vector dependent on the coupling constant, (ii) give rise to spontaneous breaking of this symmetry at $T = 0$ accompanied by generation of the magnetic moment and appearance of gapless modes with a non-relativistic (*ferromagnetic*) dispersion $E \sim k^2$, (iii) generate massive (roton) modes.

I. INTRODUCTION

Though general stability of critical points is determined by scaling dimensions of the perturbing operators, to determine the ultimate destination of the renormalization group (RG) flow is much more difficult problem. Availability of non-perturbative methods makes this task easier in (1+1)- and two dimensions.

In (1+1)- or two dimensions critical points possess conformal symmetry with the Lorentz ($O(2)$) symmetry being its part.¹ If the relevant perturbation preserves the Lorentz symmetry, the most it can do is to open a gap in a part of the spectrum. That is exactly what happens in all known solvable examples (for instance, in the sine-Gordon model).

It may happen however, that the perturbation violates the Lorentz symmetry. Such perturbations certainly make sense in condensed matter physics where preservation of the Lorentz symmetry is not required. There are quite a few models of condensed matter physics which can be treated as critical models perturbed by relevant Lorentz-symmetry-breaking operators. These operators have non-zero Lorentz spin. Since in (1+1)- or two dimensions the Lorentz symmetry at criticality is extended to conformal symmetry, Lorentz spin is also called *conformal* spin. At criticality where all two-point correlation functions follow power law behaviour, an operator $\hat{O}(z, \bar{z})$ is characterized by its scaling dimension d and conformal (or Lorentz) spin S defined as

$$\langle \hat{O}(z_1, \bar{z}_1) \hat{O}(z_2, \bar{z}_2) \rangle = A z_{12}^{-(d+S)} \bar{z}_{12}^{-(d-S)} \quad (1)$$

where $z, \bar{z}$ are holomorphic coordinates defined as $z = \tau + ix$, $\bar{z} = \tau - ix$ (in (1+1)-dimensions τ is Matsubara time). Since only bosonic operators can appear as perturbations, possible Lorentz spins are *integer*. Furthermore, since in unitary theories conformal dimensions are positive ($d \pm S \geq 0$) and $d \leq 2$ for relevant operators, this leaves us with only one choice: $S = \pm 1$.

The simplest example of $S = \pm 1$ operator is a conserved charge. At criticality such perturbation generates incommensurability. As an illustration one can consider a change of the chemical potential for massless Dirac fermions:

$$A = \int d\tau dx [R^+ \partial R + L^+ \bar{\partial} L - \mu(R^+ R + L^+ L)] \quad (2)$$

This perturbation is removed by a simple transformation of the fields

$$R \rightarrow e^{i\mu x} R, \quad L \rightarrow e^{-i\mu x} L \quad (3)$$

¹Strictly speaking, the Lorentz symmetry is violated in the models with complex isotopic symmetry, where different sectors of the spectrum may have different velocities. However, this does not influence the argument made in the text.

and thus leads to a shift in the corresponding Fermi momentum and generates static oscillations with the new wave vector whose magnitude depends on the value of μ . The perturbed theory remains critical in the infrared (IR).

The situation becomes less clear when the perturbation is not a constant of motion. Such non-trivial perturbation corresponds to a non-holomorphic operator, that is, in the notations of Eq.(1), an operator with $d \neq \pm S$.

A well studied example of $S = \pm 1$ perturbation is the the so-called Z_N 'chiral clock' model of statistical physics (see [1] for review). In fact, this model is not too good for statistical physics purposes since its Boltzmann weights are not positively defined. However, the (1+1)-dimensional version of this model makes perfect sense and, as was demonstrated in [2], describes a tunneling between two spinless Luttinger liquids. It was established that for the chiral clock model the relevant $S = \pm 1$ perturbations generate incommensurability.

An immediate generalization of the problem of tunneling includes spin. The tunneling between two spin-1/2 Luttinger liquids is described by the Hamiltonian

$$H = H_{LL}^{(1)} + H_{LL}^{(2)} + \int dx (\mathcal{T}_+ + \mathcal{T}_-), \quad (4)$$

$$\mathcal{T}_+ = t(R_{1,\sigma}^+ R_{2,\sigma} + H.c), \quad \mathcal{T}_- = t(L_{1,\sigma}^+ L_{2,\sigma} + H.c) \quad (5)$$

where $R_{1,2}$ and $L_{1,2}$ are the right- and the left-moving fermions on chains 1 and 2; H_{LL} describe interacting electrons on the individual chains. Due to the intra-chain interaction the fermionic operators acquire non-trivial scaling dimensions such that

$$\langle \mathcal{T}_\pm(\tau, x) \mathcal{T}_\pm(0, 0) \rangle \sim \frac{1}{(\tau \pm ix)^2} \frac{1}{(\tau^2 + x^2)^\theta} \quad (6)$$

where θ depends on the interaction (here I do not discuss the effects related to difference between the charge v_c and the spin v_s velocities of the excitations and set $v_c = v_s = 1$). According to definition (1) operators $\mathcal{T}_\pm$ carry conformal (Lorentz) spin $S = \pm 1$.

Another problem much discussed in the literature is the problem of frustrated spin-1/2 two-leg Heisenberg ladder (alias the zig-zag ladder). In the decoupled limit, two $S=1/2$ chains represent an $SU(2)_1 \times SU(2)_1$ WZNW theory. Each $SU(2)_1$ WZNW model has its matrix field $\hat{g}_i(x)$ ($i = 1, 2$) = $\epsilon_i(x) + \mathbf{n}_i \cdot \vec{\sigma}$, ($\mathbf{n}_{1,2}$ are staggered magnetizations of chains 1 and 2 and $\epsilon_{1,2}(x)$ are the staggered energy density operators). For the frustrated ladder the interchain interaction contains is dominated by the so-called twist term [3]. The most general $SU(2)$ -invariant form of this term is

$$\mathcal{O}_{\text{twist}} = A \mathbf{n}_1 \cdot \partial_x \mathbf{n}_2 + B \epsilon_1 \partial_x \epsilon_2 + [1 \leftrightarrow 2] \quad (7)$$

In the Heisenberg zigzag ladder the bare value of B is zero but it is generated in the course of RG (due to the fusion of the A -operator with scalar marginal ones). where the leading interchain interaction gives the following contribution to the action density [3]: For a single chain the two-point function of the operators $\mathbf{n}, \epsilon$ decays as

$$\langle \mathbf{n}(\tau, x) \mathbf{n}(0, 0) \rangle = \langle \epsilon(\tau, x) \epsilon(0, 0) \rangle \sim (x^2 + \tau^2)^{-1/2} \quad (8)$$

which means that

$$\langle \mathcal{O}(\tau, x) \mathcal{O}(0, 0) \rangle \sim (x^2 + \tau^2)^{-1} [(\tau - ix)^{-2} + (\tau + ix)^{-2}] \quad (9)$$

According to definition (1), this operator is a sum of two operators with $d = 2, S = \pm 1$.

Using the procedure suggested in [8], one can reformulate the zig-zag ladder model as a model of four Majorana fermions with the Lagrangian density given by

$$\begin{aligned} \mathcal{L} &= \mathcal{L}_0 + \mathcal{L}_{\text{twist}}, \\ \mathcal{L}_0 &= \sum_{a=0}^3 \left[\frac{1}{2} \chi_a (\partial_\tau - i \partial_x) \chi_a + \frac{1}{2} \bar{\chi}_a (\partial_\tau + i \partial_x) \bar{\chi}_a \right] \\ \mathcal{L}_{\text{twist}} &= \{ g_1 \chi_1 \chi_2 \chi_3 \bar{\chi}_0 + g_2 [\chi_0 \chi_1 \chi_2 \bar{\chi}_3 + \chi_3 \chi_0 \chi_1 \bar{\chi}_2 + \chi_2 \chi_3 \chi_0 \bar{\chi}_1] + [(\chi \leftrightarrow \bar{\chi})] \} \end{aligned} \quad (10)$$

where g_1, g_2 are linear combinations of A, B . In the limit $A = B$ we get the perfect $O(4)$ symmetry $g_1 = g_2$, with a suitable choice of A, B one can get $g_2 = 0$ which corresponds to the model I consider later in the paper.

Operators with non-zero conformal spin may also appear in models describing tunneling between edges of incompressible Fractional Quantum Hall (FQH) states [9].

Since operators with non-zero conformal spin break Lorentz symmetry, it lifts the restrictions on the low energy spectrum. With the Lorentz symmetry being preserved one has two choices: either to have a linear spectrum or to have a spectral gap. Without Lorentz symmetry one can imagine a quadratic gapless spectrum which would be naturally associated with a ferromagnetic Goldstone mode. Thus, if a model in question has a continuous isotopic symmetry, the presence of $S = \pm 1$ perturbation may lead to a spontaneously breaking of this symmetry at $T = 0$ with a formation of a net 'magnetic' moment and the above-mentioned quadratic spectrum.

In all examples discussed above (except the one involving the edge states) the perturbations with $S = \pm 1$ appear together. This introduces an additional complication because a fusion of two operators with opposite Lorentz spins generates under RG a relevant operator with zero spin. However, in a certain range of parameters of model (5) this secondary flow does not catch up with the flow of the original coupling constant (for model (7) this is always the case) [4], [5], [6] (see also [7]) where this topic is extensively discussed). This gives room for an energy scale on which the effect of interaction between the operators with $S = 1$ and $S = -1$ can be neglected. On this energy scale one can drop one operator and consider a perturbation with one sign of conformal spin, for example $S = 1$ [10]. In FQH systems, where excitations are chiral, appearance of a single $S = \pm 1$ perturbation is more likely.

II. A SOLVABLE MODEL

In the subsequent sections I discuss a series of exactly solvable models which realize the symmetry breaking scenario outlined above. These models are related to Wess-Zumino-Novikov-Witten (WZNW) model. The latter model together with the related lattice models has been used extensively to address the problems in magnetism (see, for example, [11], [12], [13], [14]). Some of these lattice models are rather similar to the zig-zag spin ladder (10).

The particular model I discuss is described by the following action:

$$A = W_k^* + h_{\bar{a}} \int d^2x : J^a(x) \Phi_{a,\bar{a}}(x) : \quad (11)$$

where W_k^* is the action of the *critical* Wess-Zumino-Novikov-Witten (WZNW) model on the $SU(2)$ group with level $k \geq 2$, $\Phi_{a,\bar{a}}$ are the primary field from the adjoint representation, J_a are the left Kac-Moody currents and h_a is a constant vector. The restriction on the value of k is related to the fact that for $k = 1$ field $\Phi_{a\bar{a}}$ does not exist. The suggested solution can be easily generalized for other symmetry groups and coset models.

To make the discussion self-contained, I recall several basic facts about WZNW models. A WZNW model describes a matrix field $g(x)$ defined on a group G whose dynamics is governed by the action

$$W(g) = \frac{1}{16\lambda\pi} \int d^2x \text{Tr}(\partial_\mu g^{-1} \partial_\mu g) + k\Gamma[g] \quad (12)$$

$$\Gamma[g] = -\frac{i}{24\pi} \int_0^\infty d\xi \int d^2x \epsilon^{\alpha\beta\gamma} \text{Tr}(g^{-1} \partial_\alpha g g^{-1} \partial_\beta g g^{-1} \partial_\gamma g)$$

The action contains two parameters - the coupling constant λ and integer number k . The model has a global $G_R \times G_L$ symmetry being invariant under the transformations

$$g(x) \rightarrow Vg(x)U \quad (13)$$

Hence there are two conserved charges associated with left and right global shifts of the matrix field g . In the perturbed model (11) the right symmetry is broken, but the left symmetry is not. As we shall see below, this symmetry is broken spontaneously at zero temperature.

Model (12) has a stable critical point; the critical value of the coupling constant is $\lambda^* = k^{-1}$. In what follows I shall distinguish between WZNW model as such and *critical* WZNW model described by action (12) with $\lambda = k^{-1}$.

At the critical point WZNW models possesses a higher symmetry: its right (left) *currents* become holomorphic (antiholomorphic):

$$\bar{J} = \frac{k}{2\pi} g^{-1} \bar{\partial} g, \quad \partial \bar{J} = 0,$$

$$J = -\frac{k}{2\pi} g \partial g^{-1}, \quad \bar{\partial} J = 0 \quad (14)$$

(where $\partial = \partial_z$, $\bar{\partial} = \partial_{\bar{z}}$; $z = \tau + ix$, $\bar{z} = \tau - ix$) and satisfy the Kac-Moody algebra:

$$[J^a(x), J^b(y)] = \frac{ik}{4\pi} \delta^{ab} \delta'(x-y) + if^{abc} J^c(y) \delta(x-y) \quad (15)$$

The critical WZNW model can be conveniently described using the Hamiltonian formalism:

$$\hat{H} = \frac{2\pi}{k + c_v} \sum_{a=1}^D \int dx [: J^a(x) J^a(x) : + : \bar{J}^a(x) \bar{J}^a(x) :]$$

where c_v is the quadratic Casimir in the adjoint representation. For the SU(2) group $c_v = 2$.

At criticality any field which is local in g can be represented as a linear combination of mutually local operators composing a basis in the operator space. This basis of fields contains the primary fields $\Phi^{(j)}$ and their descendants. The primary fields transform under $G_R \times G_L$ as tensors belonging to irreducible representations of the group; the descendants are generated from the primary fields by fusion with the Kac-Moody currents. Primary fields from higher representations can be generated by fusion of the fields from the fundamental representation. However, this process terminates after certain number of fusions depending on the value of k .

To make the discussion more concrete, I concentrate on the SU(2) group. In this case the primary fields are characterized by the value of spin $j = 1/2, 1, 3/2, \dots$. The field with the smallest spin $j = 1/2$ is the g -matrix itself. A fusion of two g -operators generates a field with $j = 1$. For the SU(2) group this representation is isomorphic to the adjoint one. The maximal spin one can achieve by fusing g -matrices is equal to $k/2$. Therefore for $k = 1$ model there is no primary field with $j = 1$ and model (11) is not well defined.

III. BETHE ANSATZ SOLUTION

To solve model (11) I use the Bethe ansatz solution of WZNW model [16]. This solution describes not just the critical point, but the entire RG flow towards it. As was shown in [17], in the vicinity of the critical point the leading irrelevant operator responsible for this flow is

$$V = \lambda : \bar{J}_a J_{\bar{a}}(x) \Phi_{a,\bar{a}}(x) : \quad (16)$$

In the presence of the right magnetic field $\mathbf{H}_R$ this operator generates perturbation (11). Indeed, this field is coupled only to the currents of right chirality $\bar{J}_{\bar{a}}$. In a finite magnetic field these currents acquire a finite expectation value

$$\langle \bar{J}_{\bar{a}} \rangle = \delta^{ab} (H_a^{(R)} / 2\pi) \quad (17)$$

and therefore in the leading order in H the irrelevant operator (16) is transformed into a relevant perturbation (11) with

$$h^a = \lambda (H_a^{(R)} / 2\pi) \quad (18)$$

Thermodynamic Bethe Ansatz (TBA) equations for SU(2) WZNW model of level k in magnetic field have the following form [16]:

$$\begin{aligned} \epsilon_n(v) &= Ts * \ln[1 + e^{\epsilon_{n+1}(v)/T}] [1 + e^{\epsilon_{n-1}(v)/T}] - m\delta_{n,0} e^{\pi v/2} - m\delta_{n,k} e^{-\pi v/2}, \\ \lim_{n \rightarrow -\infty} \frac{\epsilon_n}{n} &= -H_R, \quad \lim_{n \rightarrow \infty} \frac{\epsilon_n}{n} = H_L \end{aligned} \quad (19)$$

where $n = -\infty, \dots, \infty$. H_R and H_L are right and left 'magnetic' fields corresponding to the two conserved charges. In accordance with the above discussion, I will keep only the leading contribution in $H_R = H$ and keep H_L infinitesimal.

The free energy is given by

$$F/L = -mT \int dv (e^{\pi v/2} \delta_{n,0} + \delta_{n,k} e^{-\pi v/2}) \ln[1 + e^{\epsilon_n(v)/T}] \quad (20)$$

(L is the system's size) and

$$s * f(v) = \int_{-\infty}^{\infty} du \frac{f(u)}{4 \cosh[\pi(v-u)/2]}$$

We first study these equations at $T = 0$. In this case all ϵ_n with negative n are of order of nH , $\epsilon_0(v)$ is positive at $v \rightarrow -\infty$ and changes its sign at $B \sim -\ln(m/H)$, ϵ_n with positive n are also positive except of $\epsilon_k(v)$ which changes its sign from negative to positive at some value $Q > 0$. The ground state energy can be expressed in terms of ϵ_0 and ϵ_k which satisfy the following integral equations:

$$\int_Q^\infty du D(v-u) \epsilon_k(u) = -m e^{-\pi v/2} + \int_{-\infty}^{-B} du K(v-u) \epsilon_0(u) \quad (21)$$

$$\int_{-\infty}^{-B} du D(v-u) \epsilon_0(u) = -m e^{\pi v/2} + H + \int_Q^\infty du K(v-u) \epsilon_k(u) \quad (22)$$

where the Fourier transforms of the kernels are

$$D(\omega) = \frac{\tanh \omega e^{k|\omega|}}{2 \sinh k\omega}, \quad K(\omega) = \frac{\tanh \omega}{2 \sinh k\omega} \quad (23)$$

The limits B and Q are determined by the conditions $\epsilon_0(-B) = 0, \epsilon_k(Q) = 0$.

To determine the spectrum, we also need to know the distribution densities ρ_0, ρ_k which satisfy similar equations:

$$\int_Q^\infty du D(v-u) \rho_k(u) = \frac{m}{4} e^{-\pi v/2} + \int_{-\infty}^{-B} du K(v-u) \rho_0(u) \quad (24)$$

$$\int_{-\infty}^{-B} du D(v-u) \rho_0(u) = \frac{m}{4} e^{\pi v/2} + \int_Q^\infty du K(v-u) \rho_k(u) \quad (25)$$

These distribution densities determine the conserved charges (right and left 'magnetic' moments):

$$Q_L = \frac{1}{2} \int_Q^\infty du \rho_k(u), \quad Q_R = \frac{1}{2} \int_{-\infty}^{-B} du \rho_0(u) \quad (26)$$

Since we are interested only in the leading asymptotics in H , we can neglect ϵ_k in Eq.(22) (and respectively ρ_k in Eq.(25)) and solve these equations as Wiener-Hopf ones. Eqs.(21, 24) in this case also become Wiener-Hopf equations. The case $k = 2$ is somewhat special and will be treated separately. For $k > 2$ the solutions at large Q, B (small H/m) are given by

$$\epsilon_0(v) = \int_{-\infty}^\infty \frac{d\omega}{2\pi} e^{i\omega(v+B)} \epsilon_0^{(+)}(\omega), \quad \epsilon_0^{(+)}(\omega) = \frac{\pi H}{i\omega(-\pi + 2i\omega)G^{(+)}(\omega)G^{(+)}(0)}, \quad (27)$$

$$\epsilon_k(v) = \int_{-\infty}^\infty \frac{d\omega}{2\pi} e^{-i\omega(v-Q)} \epsilon_k^{(+)}(\omega), \quad \epsilon_k^{(+)}(\omega) = \frac{\pi(k-2)m e^{-\pi Q/2}}{(\pi - 2i\omega)(\pi - ik\omega)G^{(+)}(\omega)G^{(-)}(-i\pi/2)}, \quad (28)$$

$$\rho_k^{(+)}(\omega) = \epsilon_k^{(+)}(\omega) \frac{(\pi - ik\omega)}{2\pi(k-2)} \quad (29)$$

where

$$m \exp(-\pi B/2) = H \frac{G^{(+)}(i\pi/2)}{G^{(+)}(0)}, \quad (30)$$

$$\exp(-\pi Q/2) = \beta_k (H/m)^{\frac{(k+2)}{(k-2)}}, \quad (31)$$

$$\beta_k = \left\{ \frac{k \tan(\pi/k)}{2\pi(k+2)[G^{(+)}(i\pi/k)]^2} \right\}^{k/(k-2)} \left[\frac{G^{(+)}(i\pi/2)}{G^{(+)}(0)} \right]^{\frac{(k+2)}{(k-2)}}$$

and

$$G^{(-)}(\omega) = \left(\frac{i\omega k + 0}{\pi e} \right)^{-\frac{i\omega k}{\pi}} \frac{\Gamma(1 + \frac{i\omega k}{\pi}) \Gamma(\frac{1}{2} + \frac{i\omega}{\pi})}{\sqrt{2\pi k} \Gamma(1 + \frac{i\omega}{\pi})} \quad (32)$$

with $G^{(-)}(\omega) = G^{(+)}(-\omega)$ and $D(\omega) = G^{(-)}(\omega)G^{(+)}(\omega)$.

To obtain the modified TBA equations describing model (11) I simply replace ϵ_0 in Eq.(19) by its zero temperature approximate value (27). The result is

$$\epsilon_n(v) = Ts * \ln[1 + e^{\epsilon_{n+1}(v)/T}] [1 + e^{\epsilon_{n-1}(v)/T}] + \delta_{n,1} s * \epsilon_0(v) - m \delta_{n,k} e^{-\pi v/2}, \quad (n = 1, \dots) \quad (33)$$

$$\lim_{n \rightarrow \infty} \frac{\epsilon_n}{n} = H_L$$

The free energy is then given by

$$F/L = -mT \int dv e^{\pi v/2} \ln[1 + e^{\epsilon_k(v)/T}] \quad (34)$$

Let us return to $T = 0$ solution. From equations (28, 29) we can extract the following information. First of all, I observe that ϵ_k approaches zero at two points: at $v = Q$ and $v \rightarrow \infty$. The latter point I identify with zero momentum; then point $v = Q$ corresponds to the wave vector

$$\delta P = 2\pi \int_Q^\infty dv \rho_k(v) = 2\pi \rho_k^{(+)}(\omega = 0) = m \gamma_k (H/m)^{\frac{(k+2)}{(k-2)}} \quad (35)$$

where

$$\gamma_k = \frac{\beta_k}{G^{(+)}(i\pi/2)G^{(+)}(0)}$$

This incommensurate wave vector scales with H exactly as one expects it to scale (that is $\delta P \sim H^{1/(2-d)}$), taking into account that the scaling dimension of the perturbation in Eq.(11) is $d = 1 + 4/(k+2)$. The excitation spectrum in the vicinity of $v = Q$ is linear; the velocity is given by

$$V_R = \frac{1}{2\pi \rho_k(Q)} \frac{\partial \epsilon_k(v)}{\partial v} \Big|_{v=Q} = \frac{(k-2)}{k} V_L \quad (36)$$

where V_L is the velocity of the left-moving particles (I have been working in the system of units where $V_L = 1$).

The fact that ϵ_k approaches zero also at $v \rightarrow \infty$ means that there are soft modes at zero momentum. Since according to Eqs.(28, 29) $\epsilon_k(\omega), \rho_k(\omega)$ behave as $|\omega|$ at small ω , their real space asymptotics at $v \rightarrow \infty$ are $\epsilon_k \sim \rho_k \sim v^{-2}$. Since the momentum is given by

$$P(v) = 2\pi \int_v^\infty dv \rho_k(v) \quad (37)$$

this means that the spectrum is quadratic. Using this formula and Eq.(29) we find the dispersion law:

$$\epsilon(P) = \frac{(k-2)}{k} \frac{P^2}{\delta P}, \quad (P \ll \delta P) \quad (38)$$

The maximum of the energy is reached at $P \sim \delta P$ and is of order of δP .

For $k = 2$ the formulae for the velocity and the dispersion law should be modified:

$$Q = 2\pi/g + \frac{1}{\pi} \ln(4g), \quad g = [HG^{(+)}(i\pi/2)/m]^2 \quad (39)$$

$$V_R/V_L = \frac{1}{\pi^2} g \sim H^2, \quad (40)$$

$$\delta P = \frac{m}{H[G^{(+)}(i\pi/2)]^2} \exp\{-[\pi m/HG^{(+)}(i\pi/2)]^2\} \quad (41)$$

Taking $T \rightarrow 0$ limit in Eqs.(33) I obtain the following expressions for the energies:

$$\epsilon_n(\omega) = \frac{\sinh n\omega}{\sinh k\omega} e^{i\omega Q} \epsilon_k^{(+)}(\omega) + \frac{\sinh(k-n)\omega}{\sinh k\omega} e^{-i\omega B} \epsilon_0^{(+)}(-\omega), \quad (42)$$

$$n < k$$

$$\epsilon_{n+k}(\omega) = e^{-n|\omega|} \epsilon_k^{(+)}(\omega) e^{i\omega Q} \quad (43)$$

Let us study asymptotics of these energy functions. In real space we have

$$\epsilon_{n+k}(v) = \frac{1}{\pi} \int_Q^\infty du \frac{n}{(v-u)^2 + n^2} \epsilon_k^{(+)}(u) \quad (44)$$

From this expression it follows that $\epsilon_{n+k}(v)$ decay as v^{-2} on both infinities. On the other hand, $\epsilon_n(v)$ with $n < k$ decay as v^{-2} only at $v \gg Q$. At $v < Q$ they have roton-like minima. For $Q - v \gg 1$ I obtain from Eq.(42):

$$\begin{aligned} \epsilon_n(v) &\approx k^{-1} \sin(\pi n/k) \left[\epsilon_k^+(i\pi/k) e^{\pi(v-Q)/k} + \epsilon_0^+(i\pi/k) e^{-\pi(v+B)/k} \right] \\ &= 2k^{-1} \sin(\pi n/k) [\epsilon_k^+(i\pi/k) \epsilon_0^+(i\pi/k)]^{1/2} e^{-\pi(Q+B)/2k} \cosh[\pi(v-v_0)/k] \end{aligned} \quad (45)$$

which corresponds to the relativistic spectrum with spectral gaps for $k \neq 2$ ($n = 1, \dots, k-1$) given by

$$M_n = M_0 \sin(\pi n/k), \quad M_0 = \left[\frac{2(k-2) \cot(\pi/k)}{\pi k(k+2)} \right]^{1/2} \delta P \quad (46)$$

For $k = 2$ the gap is exponential in H^{-2} . The energy minimum for ϵ_n with $n = 1, \dots, k-1$ occurs at

$$v_0 = Q - \frac{k}{2\pi} \ln \left\{ \frac{(k-2) \tan(\pi/k)}{4\pi(k+2)[G^{(+)}(i\pi/k)]^2} \right\} \quad (47)$$

This means two things. The first one is that the massive modes are centered at *incommensurate* wave vector. The second is that in the rapidity space they are very close to Q .

IV. BETHE ANSATZ DERIVATION OF THE LOW ENERGY EFFECTIVE ACTION

Thus we have the following regions in rapidity space where low-energy excitations are located:

- (i) at $v \gg Q$ there are gapless modes ϵ_n ($n = 1, \dots$) with the spectrum $\sim v^{-2}$;
- (ii) at $v \ll Q$ there are gapless modes ϵ_n ($n = k+1, \dots$) with the spectrum $\sim v^{-2}$;
- (iii) at $v = Q$ there is a gapless mode ϵ_k with the spectrum $\sim (v-Q)$;
- (iv) at $v \approx Q$ there are massive modes ϵ_n ($n = 1, \dots, k-1$). The massive modes can be treated as low energy excitations only for $k \gg 1$ when their masses are much smaller than δP .

In the momentum space the spectrum in sectors (i),(ii) is quadratic $\epsilon(P) \sim P^2$ (see Figs. 1,2), the spectrum in sector (iii) is linear $\epsilon(P) \sim |P - P(Q)|$ and the spectrum in sector (iv) is massive relativistic (see Fig.3).




FIG. 1. Dispersion $\epsilon_n(P)$ ($n = k+1, \dots$)




FIG. 2. Dispersion $\epsilon_n(P)$ ($n = 1, \dots, k-1$)




FIG. 3. Dispersion $\epsilon_k(P)$

Let us first consider the case of moderate k when one does not need to consider roton modes. Then the low energy sector is described by the truly gapless modes (i - iii). Since their positions in rapidity space are well separated from each other and the integration kernels in Eqs.(33) decay exponentially, one can derive separate sets of TBA equations for each mode. Comparing these equations with TBA equations for known integrable models one can deduce the effective action for the low energy sector.

Let us derive TBA equations for sector (i). In this sector all energies are positive. Therefore it is convenient to have TBA equations in such a form where the integral kernels act on functions $\ln(1 + e^{-\epsilon_n/T})$. Such equations would provide a ready $T = 0$ limit. To get such form of TBA. The resulting equations read

$$T \ln[1 + e^{\epsilon_n(v)/T}] - A_{nm} * T \ln[1 + e^{-\epsilon_m(v)/T}] = A_{k,k}^{-1} * A_{n,k} * \epsilon_k^{(0)}(v) + nH_L, \quad n = 1, \dots, \quad v \gg Q \quad (48)$$

where $\epsilon_k^{(0)}(v)$ is the solution at $T = 0$ and is given by Eq.(28). The kernels have the following standard Fourier transforms:

$$A_{nm}(\omega) = \coth |\omega| [\exp(-|n-m||\omega|) - \exp(-|n+m||\omega|)] \quad (49)$$

These equations are valid for temperatures $T \ll \delta P$ - the maximum value of $\epsilon_k^{(0)}(v)$. In this region These equations coincide with the *right chiral* sector of the spin-($k/2$) integrable *ferromagnet*. The latter ones can be extracted from TBA equations obtained in [18]. In a similar fashion I obtain TBA for sector (ii):

$$T \ln[1 + e^{\epsilon_{n+k}(v)/T}] - A_{nm} * T \ln[1 + e^{-\epsilon_{m+k}(v)/T}] = a_n * \epsilon_k^{(0)} + nH_L, \quad n = 1, \dots, \quad v \ll Q \quad (50)$$

These equations describe the *left chiral* sector of the spin-1/2 integrable *ferromagnet* where the dispersion law is $\epsilon(k) \sim k^2 \theta(k)$. Therefore the parity in the ferromagnetic sector is broken.

Another soft mode in the right chiral sector has the linear spectrum with velocity (36) and is centered at the momentum δP . Using the standard manipulations with TBA equations one can find its contribution to the specific heat and establish that this mode carries central charge 1. Therefore it is described by the *non-chiral* Gaussian model.

V. CONNECTION TO THE ZIG-ZAG CHAIN AT $K = 2$

For $k = 2$ model (11) simplifies considerably. In this case the critical WZNW action is equivalent to the theory of three massless Majorana fermions $\chi_a, \bar{\chi}_a$ ($a = 1, 2, 3$) and the adjoint operator and the currents are given by [15]

$$\Phi_{ab} = \chi_a \bar{\chi}_b, \quad J^a = \frac{i}{2} \epsilon^{abc} \chi_b \chi_c \quad (51)$$

Therefore for $k = 2$ one can rewrite the perturbed action (11) solely in terms of Majorana fermions:

$$A = \int d^2x \left[\frac{1}{2} \chi_a (\partial_\tau - i\partial_x) \chi_a + \frac{1}{2} \bar{\chi}_a (\partial_\tau + i\partial_x) \bar{\chi}_a + h_a \chi_1 \chi_2 \chi_3 \bar{\chi}_a \right] \quad (52)$$

In this form the model closely resembles the model for the zig-zag ladder (10) (see also [10]) and at $g_2 = 0$ it even coincides with model (11) exactly solvable by the Bethe ansatz.

Acknowledgements I am grateful to Ph. Lecheminant for inspiration, to A. A. Nersesyan, D. Maslov and F. H. L. Essler for illuminating discussions and interest to the work and E. Papa and S. Carr for help with preparation of the manuscript.

- [1] M. den Nijs, in *Phase Transitions and Critical Phenomena*, ed. by C. Domb and J. Lebowitz, vol. 12, (Academic Press, 1988).
- [2] P. Fendley and C. Nayak, cond-mat/0008157.
- [3] A. A. Nersesyan, A. O. Gogolin and F. H. L. Essler, Phys. Rev. Lett. **81**, 910 (1998).
- [4] F. V. Kusmartsev, A. Luther and A. A. Nersesyan, JETP Lett. **55**, 692 (1992); Phys. Lett. **A176**, 363 (1993).
- [5] V. M. Yakovenko, JETP Lett. **56**, 5101 (1992).
- [6] D. V. Khveshchenko and T. M. Rice, Phys. Rev. **B50**, 252 (1993); D. V. Khveshchenko, Phys. Rev. **B50**, 380 (1993).
- [7] For references and discussion see A. O. Gogolin, A. A. Nersesyan and A. M. Tsvelik, 'Bosonization and Strongly Correlated Systems', Chapter 8, 20; Cambridge University Press (1998).
- [8] D. G. Shelton, A. A. Nersesyan and A. M. Tsvelik, Phys. Rev. B (1996).
- [9] I am grateful to S. L. Sondhi for bringing this possibility to my attention.
- [10] M. Zarea, M. Fabrizio and A. A. Nersesyan, in preparation.
- [11] L.D.Faddeev and N. Yu. Reshetikhin, Ann.Phys.NY **167** 227 (1986).
- [12] H.J.de Vega and F.Woynarovich, J.Phys.A, **25** 4499 (1992)
- [13] P.Schlottmann, Phys.Rev.B, **49**, 9202 (1994); A. A. Zvyagin and P.Schlottmann, ibid., **52** 6569 (1995).
- [14] A. A. Zvyagin, Low Temp.Phys. **26** 134 (2000) (see also cond-mat/0002356).
- [15] A. B. Zamolodchikov and V. A. Fateev, Yad. Fiz. [Sov. J. Nucl. Phys.] **43**, 1043 (1986).
- [16] A. M. Polyakov and P. B. Wiegmann, Phys. Lett. **B131**, 121 (1983); **141**, 223 (1984).
- [17] V. G. Knizhnik and A. B. Zamolodchikov, Nucl. Phys. **B247**, 83 (1984).
- [18] H. M. Babujan, Nucl. Phys. **B215**, 317 (1983).
- [19] A. M. Tsvelick, Nucl. Phys. **B305**, 675 (1988).
- [20] E. Lieb and D. Mattis, Phys. Rev. **125**, 164 (1962)